



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

average height, with shorter heads and fewer grains, thus emphasizing the importance of selecting the superior plants instead of the superior individual grains. As this variety of oats was undoubtedly a mixture of several distinct biotypes, it does not follow that the same mathematical results would be found in other varieties composed of different mixtures. The variety with which WALDRON worked may have contained a short-headed, short-culmed, heavy-grained biotype. In some other mixture the heavy-grained biotype might have longer culms, longer heads, and more numerous grains, and it would then give a positive correlation where WALDRON found a negative correlation, but this does not lessen the importance of the conclusion reached that the individual plant and not the individual grain is the proper unit of selection.—GEO. H. SHULL.

Anatomy of the seedling of *Trapa*.—A short paper by QUEVA¹¹ on the curious seedling of *Trapa natans* recalls the case of "caulicle" vs. "radicle," the question of the importance of the root as a primarily essential part of a seed plant. The author confirms the observations of previous investigators concerning the marked inequality of the cotyledons, the negative geotropism of the caulicle (which he prefers to call the hypocotyledonary axis), and the presence of internal phloem in the stem and leaves. His own investigation has resulted in the discovery of this internal phloem in the hypocotyledonary axis and in the petiole of the cotyledon. Although he finds in the very tip of the hypocotyledonary axis a vascular condition which is peculiarly root-like (two xylem points alternating with two small groups of phloem), yet he thinks it is not root, because (1) there is no rotation of the strands in successive levels from the tip of the organ to the cotyledonary node; (2) the xylem points are too near the periphery of the cylinder to look like root poles; and (3) the whole organ is covered by epidermis, except at the spot where the suspensor was attached. The growth of the caulicle, or hypocotyledonary axis, is limited; the roots strike out from its side, their vascular strands being inserted on certain metaxylem elements discernible in cross-section on one side of the vascular cylinder.—SISTER HELEN ANGELA.

Sporophylls of *Selaginella*.—SYKES and STILES¹² have made a very interesting study of the sporophylls of *Selaginella*, finding an amount of variation and a degree of complexity that have not attracted attention heretofore. A few of the more representative species are described and the different forms of the sporophyll are pointed out as "special adaptations for the secure protection of the sporangia." In many sporophylls there is a well developed air cavity in the base, and the authors suggest "that they recall the mucilage

¹¹ QUEVA, C., Observations anatomiques sur le *Trapa natans* L. Compt. Rend. Assoc. Fran. Av. Sci. 1909. Congrès de Lille, pp. 512-517. figs. 2. 1909.

¹² SYKES, M. G., and STILES, W., The cone of the genus *Selaginella*. Annals of Botany 24:523-536. pl. 41. 1910.